arXiv:cond-mat/9806256v1 22 Jun 1998

Ground-state energy of confined charged bosons in two dimensions

A. Gonzalez[1,2*], B. Partoens[3∘], A. Matulis[3◁] and F. M. Peeters[3⋄]
[1] *Departamento de Fisica, Universidad de Antioquia, AA 1226, Medellin, Colombia*
[2] *Instituto de Cibernetica, Matematica y Fisica Calle E 309, Vedado, Habana 4, Cuba*
[3] *Departement Natuurkunde, Universiteit Antwerpen (UIA), Universiteitsplein 1, B - 2610 Antwerpen, Belgium*

(February 1, 2008)

Abstract

The Padé approximant technique and the variational Monte Carlo method are applied to determine the ground-state energy of a finite number of charged bosons in two dimensions confined by a parabolic trap. The particles interact repulsively through a Coulombic, $1/r$, potential. Analytic expressions for the ground-state energy are obtained. The convergence of the Padé sequence and comparison with the Monte Carlo results show that the error of the Padé estimate is less than 4 % at any boson density and is exact in the extreme situations of very dilute and high density.

Typeset using REVTEX

The interest in finite boson systems which are confined in traps increased suddenly after the experimental observation of signals of Bose-Einstein (BE) condensation in clusters of alkali-vapour atoms.[1] These are confined three-dimensional dilute systems interacting through short-range potentials, which basically can be described by a modification of Bogoliubov's theory.[2]

Lower-dimensional systems as well as bosons interacting through long-range potentials are also widely studied. These studies are motivated, for example, by proposals of obtaining quasi two-dimensional clusters by making the traps highly anisotropic.[3] Theoretically some regularities of the spectra of energy levels have been identified to check experimentally the effective quasi-two-dimensionality of the system.[4] A particularly interesting aspect of trapped systems is related to the fact that quantum fluctuations do not destroy BE condensation in lower dimensions.[5] On the other hand, the ground-state properties of the homogeneous charged Bose gas has been studied in 3D.[6] Vortex systems in high-T_c superconductors have been shown to be equivalent to 2D bosons with logarithmic interactions.[7]

In the present paper, we study a model of two-dimensional charged bosons confined in a parabolic trap. Instead of an often used long-wavelength field theory description, we start from the N-particle hamiltonian, which in harmonic oscillator units (i.e. $\sqrt{\hbar/(m\omega_0)}$ for length, $\hbar\omega_0$ for energy, etc.) reads

$$\frac{H}{\hbar\omega_0} = \frac{1}{2}\sum_{i=1}^{N}(\vec{p}_i^{\,2} + \vec{r}_i^{\,2}) + \beta^3 \sum_{i<j} \frac{1}{|\vec{r}_i - \vec{r}_j|}, \tag{1}$$

where ω_0 is the dot confinement frequency, $\beta^3 = \sqrt{(\frac{mq^4}{\hbar^2})/(\hbar\omega_0)}$, and m and q are, respectively, the particle mass and charge. By varying β we modify the "density" of the system.

Following our approach for electrons in a parabolic quantum dot,[8] we will construct two-point Padé approximants for the ground-state energy of the Hamiltonian (1) from the weak and strong interaction expansions of the energy

$$\epsilon|_{\beta\to 0} = b_0 + b_3\beta^3 + b_6\beta^6 + \ldots, \tag{2}$$

$$\epsilon|_{\beta\to\infty} = \beta^2\{a_0 + a_2/\beta^2 + \ldots\}. \tag{3}$$

When $\beta << 1$, we have a system of weakly interacting bosons in a harmonic potential. For $\beta = 0$ we have N non-interacting bosons with energy $b_0 = N$ which is the leading contribution to the energy (2). Perturbation theory is applied to compute b_3 and b_6, yielding

$$b_3 = <0|V|0> = \frac{N(N-1)}{2}\sqrt{\frac{\pi}{2}}, \tag{4}$$

$$\begin{aligned} b_6 &= \sum_{int} \frac{<0|V|int>^2}{\epsilon_0(0) - \epsilon_0(int)} \\ &= c_1\, N(N-1)^2 + c_2\, N(N-1), \qquad (5) \\ c_1 &= -0.05300695, \quad c_2 = -0.11982050, \end{aligned}$$

where $V = \sum_{i<j} |\vec{r}_i - \vec{r}_j|^{-1}$, and int is a shorthand notation for the harmonic oscillator intermediate states respecting conservation of angular momentum (equal to zero in the ground state). Due to the pair character of the potential, only one- and two-particle excitations

are allowed. $\epsilon_0(int)$ is the energy of the non-interacting intermediate state and $\epsilon_0(0) = N$. We would like to notice that in order to apply the Padé approximant technique one needs to calculate the coefficients with rather high accuracy. This was achieved in the following way. The one-particle excitation part (namely the coefficient c_1) was calculated from the straightforward summation of a single sum which converges rather fast. The convergence of the double sum in the two-particle excitation part, however, is slow. Therefore we used an alternative approach and calculated the coefficient c_2 from the solution of the two particle interaction problem which is given in the Appendix.

On the other hand, when $\beta \to \infty$, a scaling of coordinates $r \to \beta r$ in the Hamiltonian (1) shows[8−10] that the potential energy (Coulomb repulsion plus parabolic confinement) behaves as β^2, whereas the kinetic energy ("fluctuations") is of order zero in β. Consequently, in this limit the system behaves as N classical point particles and a_0 is the minimum of the classical potential energy. The next term a_2 takes account of the zero-point fluctuations, which is given by $a_2 = \sum_\alpha \omega_\alpha/2$. The coefficients a_0 and a_2 for $N = 2 - 5$ are given in Ref. 10. They were obtained from a strong-coupling expansion of the exact quantum-mechanical Hamiltonian. For larger N, a_0 and a_2 were obtained by minimising the potential energy with a combination of Monte Carlo and Newton methods and the normal frequencies were computed by solving a classical small-oscillation problem.[11,12] Fitted expressions (as given in Ref. 8) for the coefficients a_0 and a_2 for $6 \leq N \leq 210$ are

$$a_0/N^{5/3} = 1.062 - 0.875/N^{1/2} - 0.185/N, \tag{6}$$

$$a_2/N^{5/4} = 0.573 + 0.475/N^{1/2} - 0.160/N. \tag{7}$$

These expressions also have the correct $N \to \infty$ asymptotic behavior.

Two-point Padé approximants $\{P_{s,t}(\beta)\}$ are used to estimate the ground-state energy. They are quotients of polynomials reproducing the first $s + 1$ terms of the expansion (2) and the first $t + 1$ terms of the $\beta \to \infty$ series (3). By construction, the approximants are asymptotically exact in both the $\beta \to 0$ and $\beta \to \infty$ limits. As it is common with the Padé approximants, we shall check the convergence of a sequence $\{P_{s,t}(\beta)\}$ running parallel to the diagonal $\{P_{s,s}(\beta)\}$.[13] In the present problem we use the off-diagonal sequence $\{P_{K+3,K}(\beta)\}$, which is the first non-trivial sequence with no singularities. We will show that it exhibits good convergence properties at intermediate β values. The explicit form of the first elements of this sequence are the following

$$P_{3,0}(\beta) = b_0 + \frac{b_3\beta^3}{1 + b_3\beta/a_0}, \tag{8}$$

$$P_{4,1}(\beta) = b_0 + b_3\beta^3 \left\{1 - \frac{q_1^2\beta^2}{1 + q_1\beta + q_1^2\beta^2}\right\}, \tag{9}$$

$$q_1 = b_3/a_0,$$

$$P_{5,2}(\beta) = b_0 + b_3\beta^3 \left\{1 - \frac{q_3\beta^3}{1 + q_1\beta + q_2\beta^2 + q_3\beta^3}\right\}, \tag{10}$$

$$q_1 = a_0q_2/b_3, \; q_3 = b_3q_2/a_0,$$

$$q_2 = \left(\frac{a_2 - b_0}{a_0} + \frac{a_0^2}{b_3^2}\right)^{-1},$$

$$
\begin{aligned}
P_{6,3}(\beta) = b_0 + &\frac{b_6\beta^6}{1+q_1\beta+\ldots+q_4\beta^4} \\
&+b_3\beta^3\left\{1-\frac{q_4\beta^4}{1+q_1\beta+\ldots+q_4\beta^4}\right\}, \\
q_2 &= a_0q_3/b_3,\ \ q_4=(b_3q_3+b_6)/a_0, \\
q_1 &= (q_3(b_0-a_2)+b_3)/a_0, \\
q_3 &= \frac{(a_2-b_0)|b_6|/a_0+b_3^2/a_0}{a_0^2/b_3+2b_3(a_2-b_0)/a_0}.
\end{aligned}
\tag{11}
$$

In Figs. 1(a) and 1(b) we show the relative differences between consecutive approximants for 20 and 210 bosons, respectively. From these figures, the maximum error of the $P_{6,3}(\beta)$ approximant is estimated to be lower than 4% for any value of β. For $N = 20(210)$ the maximum error is 3(4)% which is reached for $\beta = 0.8(0.4)$.

The explicit form of the coefficients a_k and b_k suggests that ϵ/N is an "almost universal" function of the variable $N\beta^3$ for large N. We found that for $N \geq 90$, the coefficients reach their asymptotic forms, and we have

$$b_k\beta^k \approx N\tilde{b}_k(N\beta^3)^{k/3}, \tag{12}$$

$$a_k/\beta^k \approx N^{5/3}\tilde{a}_k/(N\beta^3)^{k/3}, \tag{13}$$

where $\tilde{b}_0 = 1$, $\tilde{b}_3 = \frac{1}{2}\sqrt{\frac{\pi}{2}}$, $\tilde{b}_6 = -0.05300695$, $\tilde{a}_0 = 1.062$ and $\tilde{a}_2 = 0.573N^{1/4}$. The coefficients $\tilde{b}_k$ and $\tilde{a}_k$ are numbers, except for the coefficient $\tilde{a}_2$ which exhibits a smooth dependence on N. In the interval $90 \leq N \leq 210$ $\tilde{a}_2$ is approximately a constant (≈ 2) and consequently for the energy we obtain

$$\epsilon(\beta) \approx Nf(N\beta^3), \tag{14}$$

where f is only a function of $N\beta^3$. The $P_{6,3}(\beta)$ estimates for $N = 20, N = 90$ and $N = 210$ are given in Fig. 2, showing the "approximate" scaling when $N \geq 90$. Note that bosons interacting through short-range potentials also show scaling behaviour in the large-N limit. Indeed, in this limit they are described by the Gross-Pitaevskii equation, which may be written in a scaled form.[14]

In order to have an independent check on the accuracy of the present Padé approximant, we carried out variational Monte Carlo (VMC) calculations[15] for the ground-state energy. The trial wave function used in the computations was (up to a normalisation constant)

$$\Psi_T = \left(\prod_{i=1}^{N}\phi(r_i)\right)e^{-\sum_{i<j}u(r_{ij})}, \tag{15}$$

where $\phi(r) = e^{-r^2/2}$ is the ground-state function of a boson in the harmonic potential, and the pseudopotential $u(r_{ij})$ was chosen as

$$u(r) = \frac{\beta^3}{1+r^3}(A - r - \frac{C}{1+\beta}r^3\ln r), \tag{16}$$

where A and C are variational parameters. This interpolative expression for u gives the correct asymptotic form at coincidence, $r \to 0$ ("cusp conditions"),

$$u(r)|_{r\to 0} = \text{const} - \beta^3 r + \ldots, \tag{17}$$

and at very large values of r,

$$u(r)|_{r\to\infty} = -C\frac{\beta^3}{1+\beta}\ln\, r + \ldots. \tag{18}$$

The later expression can be easily verified for the $N = 2$ system. The coefficient in front of $-\ln\, r$ is roughly the difference $\epsilon(\beta) - \epsilon(0)$, which is proportional to β^3 for small β values, and tends to β^2 as β increases.

Fig. 3 compares the Monte Carlo and $P_{6,3}(\beta)$ Padé results for 20 and 210 bosons as function of β in the interval $0 \leq \beta \leq 1.3$. In the computations, 10^5 sweeps were used to compute mean values after 2×10^4 steps for thermalisation. Note that $\beta = 1.3$ is well outside the perturbative regime. Indeed, $\epsilon(1.3)/\epsilon(0) > 12$ for $N = 20$, whereas $\epsilon(1.3)/\epsilon(0) > 35$ for $N = 210$.

In Fig. 1 we have also plotted (the dots) the relative difference between the $P_{6,3}(\beta)$ Padé approximant and the variational Monte Carlo results. Note that in Fig. 1(a) for $N = 20$ up to $\beta \simeq 2.5$ (the solid dots) the $P_{6,3}(\beta)$ approximant gives practically identical results as those obtained from the variational Monte Carlo. The relative error is less than 1.5% which is below the 4% level estimated previously from a comparison between successive Padé approximants. For larger values of β one approaches the Wigner limit where the particles form a crystal-like structure. In that limit the functional form taken for the trial function is expected to be no longer good and we notice that the relative error does not decrease with β (open dots) but stays approximately constant. Also for $N = 210$ it is shown in Fig. 1(b) that the relative error is less than 4%.

ACKNOWLEDGMENTS

AG acknowledges support from the Colombian Institute for Science and Technology (COLCIENCIAS). BP is an aspirant, and FMP a research director with the Flemish Science Foundation (FWO Vlaanderen). Part of this work is supported by FWO, a NATO Linkage grant, the ‘Interuniversity Poles of Attraction Programme - Belgian State, Prime Minister’s Office - Federal Office for Scientific, Technical and Cultural Affairs’, and the WOG on field theory and statistical physics.

REFERENCES

[*] Electronic mail: agonzale@fisica.udea.edu.co

[°] Electronic mail: bpartoen@uia.ua.ac.be

[◁] Permanent address: Semiconductor Physics Institute, Gostauto 11, 2600 Vilnius, Lithuania. Electronic mail: matulis@pub.osf.lt

[⋄] Electronic mail: peeters@uia.ua.ac.be

[1] M. H. Anderson, J. R. Ensher, M. R. Mathews, C. E. Wieman, and E. A. Cornell, Science **269**, 198 (1995); K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Phys. Rev. Lett. **75**, 3969 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).

[2] See, for example, S. Giorgini, L. P. Pitaevskii and S. Stringari, J. Low Temp. Phys. **109**, 309 (1997), and references therein.

[3] M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. M. Kurn, D. S. Durfee, C. G. Townsend, and W. Ketterle, Phys. Rev. Lett. **77**, 988 (1996); C. J. Myatt, E. A. Burt, R. W. Ghrist, E. A. Cornell, and C. E. Wieman, Phys. Rev. Lett. **78**, 586 (1997).

[4] L. P. Pitaevskii and A. Rosch, Phys. Rev. A **55**, 835 (1997).

[5] Tin-Lun Ho and Michael Ma, cond-mat/9703034.

[6] V. Apaja, J. Halinen, V. Halonen, E. Krotscheck, and M. Saarela, Phys. Rev. **B 55**, 12925 (1997); G. Sugiyama, C. Bowen, and B. Alder, Phys. Rev. **B 46**, 13042 (1992); S. Moroni, S. Conti, and M. P. Tosi, Phys. Rev. **B 53**, 9688 (1996); K. Tankeshwar, B. Tanatar, and M. P. Tosi, ICTP preprint IC/97/95.

[7] D. R. Nelson, Phys. Rev. Lett. **60**, 1973 (1989); D. R. Nelson and H. S. Seung, Phys. Rev. **B 39**, 9153 (1989); W. R. Magro and D. M. Ceperley, Phys. Rev. Lett. **73**, 826 (1994); H. Nordborg and G. Blatter, Phys. Rev. Lett. **79**, 1925 (1997).

[8] A. Gonzalez, B. Partoens, and F. M. Peeters, Phys. Rev. **B 56**, 15740 (1997).

[9] A. Matulis and F. M. Peeters, J. Phys.: Condens. Matter **6**, 7751 (1994).

[10] A. Gonzalez, J. Phys.: Condens. Matter **9**, 4643 (1997).

[11] V. M. Bedanov and F. M. Peeters, Phys. Rev. **B 49**, 2667 (1994).

[12] V. A. Schweigert and F. M. Peeters, Phys. Rev. **B 51**, 7700 (1995).

[13] A. H. MacDonald and D. R. Ritchie, Phys. Rev. **B 33**, 8336 (1986).

[14] S. Giorgini, L. P. Pitaevskii, and S. Stringari, Phys. Rev. Lett. **78**, 3987 (1997).

[15] D. M. Ceperley, *Lectures on quantum Monte Carlo*, in Spring Colleges in Computational Physics, ICTP, Trieste, 1997.

FIGURES

FIG. 1. The relative differences between consecutive Padé approximants of the sequence $P_{K+3,K}(\beta)$. We consider two systems: (a) $N = 20$ and (b) $N = 210$. The dots give the relative difference $|E_{VMC} - P_{6,3}(\beta)|/P_{6,3}(\beta)$ between the best Padé approximant and the variational Monte Carlo (VMC) results.

FIG. 2. Approximate scaling properties of the energy for $N = 90$ and $N = 210$. As a comparison we also show the result for $N = 20$. The dots (solid dots for $N = 210$, open dots for $N = 90$ and stars for $N = 20$) are the results from the variational Monte Carlo calculation.

FIG. 3. Comparison between the $P_{6,3}(\beta)$ Padé estimates (solid curves) and the variational MC calculations (dashed curves) for: (a) $N = 20$ and (b) $N = 210$.

APPENDIX: SECOND ORDER ENERGY CORRECTION FOR THE TWO PARTICLE INTERACTION PROBLEM

Here we present an alternative but more accurate calculation of the coefficient c_2 for the second order energy correction (5). Because expression (5) exhibits a rather simple dependence on the number of bosons N we can limit ourselves to calculate the coefficient c_2 for the system of two particles with Coulomb repulsion in a parabolic confinement potential. The ground-state of that system will be symmetric with respect to the permutation of the particles what automatically takes into account the boson character of the problem.

It is known that in this two particle interaction problem the relative and the center-of-mass motion can be separated. The center-of-mass ($\vec{R} = (\vec{r}_1 + \vec{r}_2)/2$) motion is the one of a harmonic oscillator and is easily eliminated. Consequently, we have only to consider the following radial Schrödinger equation for the relative motion ($\vec{r} = \vec{r}_1 - \vec{r}_2$) of particles

$$\{H_0 + \lambda/r - E\}\, R(r) = 0, \tag{A1}$$

$$H_0 = -\frac{1}{r}\frac{d}{dr}r\frac{d}{dr} + \frac{1}{4}r^2, \tag{A2}$$

with $\lambda = \beta^3$.

Now let us expand the eigenvalue $E = E_0 + \lambda E_1 + \cdots$ and the wave function $R = R_0 + \lambda R_1 + \cdots$ into powers of λ. The application of the standard perturbation technique leads to the following expressions

$$E_0 = 1, \tag{A3}$$

$$E_1 = \int_0^\infty dr R_0^2(r) = \sqrt{\pi/2}, \tag{A4}$$

$$E_2 = \int_0^\infty dr R_0(r) R_1(r), \tag{A5}$$

with $R_0 = \exp(-r^2/4)$ and the first order wave function correction obeys the following equation

$$\{H_0 - E_0\}\, R_1 = (1/r - E_1)R_0. \tag{A6}$$

This equation has to be solved together with the boundary and orthogonality conditions

$$R_1(0) = R_1(\infty) = 0, \qquad \int_0^\infty drrR_1(r)R_0(r) = 0. \tag{A7}$$

Inserting the substitution $R_1 = \{\sum_{n=1}^{\infty} w_n r^n + C\}R_0$ into equation (A6) we obtain the recurrence relation

$$w_{n+2} = w_n n/(n+2)^2, \quad w_1 = 1, \quad w_2 = -E_1/4. \tag{A8}$$

Now iterating the above relation, withy the ortogonality condition (A7), and inserting the obtained result into expression (A5) we obtain the second order energy correction expressed as a single sum of Γ-functions

$$\begin{aligned} E_2 = {} & \sqrt{\frac{\pi}{16}} \sum_{k=0}^{\infty} \left\{ \frac{\Gamma(k+1)}{(k+1/2)\Gamma(k+3/2)} - \frac{\Gamma(k+3/2)}{(k+1)\Gamma(k+2)} \right\} \\ & - \frac{\pi}{8} \sum_{k=0}^{\infty} \frac{1}{(k+1/2)(k+1)}. \end{aligned} \tag{A9}$$

The coefficient $c_2 = E_2/2 - c_1$ follows from expression (5) with $N = 2$. The sums in expression (A9) converge which enables us to obtain the coefficient c_2 up to the desired accuracy.

Figure 1


(a)
N=20
K=0
K=1
K=2
0.00
0.05
0.10
0.15
0.20
0.0
0.2
0.4
0.6
0.8
1.0
$|P_{K+4,K+1}(\beta)-P_{K+3,K}(\beta)|/P_{K+4,K+1}(\beta)$
(b)
N=210
K=0
K=1
K=2
0.00
0.05
0.10
0.15
0.20
0.0
0.2
0.4
0.6
0.8
1.0
$|P_{K+4,K+1}(\beta)-P_{K+3,K}(\beta)|/P_{K+4,K+1}(\beta)$
$\beta/(\beta+1)$

Figure 2



Figure 3


N=20
Variational Monte Carlo
$P_{6,3}(\beta)$
Energy
200
150
100
50
(a)
0.0
0.2
0.4
0.6
0.8
1.0
1.2
N=210
Energy
12000
10000
8000
6000
4000
2000
0
(b)
0.0
0.2
0.4
0.6
0.8
1.0
1.2
β